2020

More than **3.6 million** U.S. youth are using e-cigarettes

How much do you know about this crisis?

Youth use of e-cigarettes, also known as vaping, remains a public health crisis. E-cigarettes are the most commonly used tobacco product among both middle and high school students.⁹ Compared to 2019, 1.8 million fewer U.S. youth are currently using e-cigarettes. However, due to alarming increases since 2011, the number of current youth e-cigarette users remains concerningly high.¹

SOME TEENS REPORT USING E-CIGARETTES IN SCHOOL BATHROOMS AND EVEN IN THE CLASSROOM.²

Learning more about the different types of e-cigarette products is an important first step in addressing youth vaping.

DID YOU KNOW:

E-cigarettes come in a variety of shapes and sizes and may not look like a tobacco product, which can make them hard to spot.²

Some devices popular among teens — like Juul and Puff Bar — are as small as a USB flash drive and even look like one.^{2,16}

Certain products emit very low amounts of aerosol or "vapor," which makes them easier to use discreetly than combustible cigarettes.^{9,11}

Most e-cigarettes contain nicotine, the same highly addictive drug in cigarettes. Some e-cigarettes⁵ may contain as much nicotine as a pack of 20 regular cigarettes.³ A Small Device... A BIG PROBLEM In 2020, approximately I in 5 high school students and 1 in 20 middle school students

CENTER FOR TOBACCO PRODUCTS

Source: Wang, et al. MMWR 2020 Note: All numbers presented here are estimates.

www.fda.gov/tobacco



f facebook.com/fda

Many teens have dangerous misperceptions that lead them to believe that vaping is harmless.

Important facts to share with youth

Vape aerosol can contain harmful chemicals

Vaping can expose the user's lungs to harmful chemicals like formaldehyde, acrolein, and acetaldehyde, which are known to cause irreversible lung damage.⁴⁵

There can be danger behind the flavor

Vapes get their flavors from chemicals. While these flavorings are safe to eat in food, they're not safe to inhale. Inhaling flavor chemicals can harm your lungs.⁶ Most vapes contain nicotine, which is highly addictive

Vaping delivers nicotine to the brain in as little as 10 seconds.⁷ A teen's brain is still developing, making it more vulnerable to nicotine addiction.^{8,14} Nicotine exposure during the teen years can disrupt normal brain development.^{9,14} It may have long-lasting effects, like increased impulsivity and mood disorders.^{9,12,13,15} Vapers could be inhaling metal particles into their lungs

Vape aerosol could be delivering metal particles like chromium, nickel, lead, tin and aluminum right into your lungs. Some of these metals are toxic.^{10,11}

FDA's Efforts to Curb Youth E-Cigarette Use

FDA is committed to protecting youth from the dangers of e-cigarettes, including cracking down on illegal sales to anyone under 21 and holding retailers and manufacturers accountable for marketing practices. Also, in addition to our national peer-to-peer public education campaign called "The Real Cost" FDA has joined forces with Scholastic to provide teachers and school administrators with the resources they need to educate their students about e-cigarettes.

Together, we've created **free lesson plans**, **activities and videos** for teachers to educate their students on the health risks of e-cigarette use. Please visit the <u>Scholastic youth</u> <u>vaping risks site</u> to access these resources in English and Spanish.

Quitting Help Is Available

There is an urgent need to share quitting resources with teens who are addicted to e-cigarettes. If you identify teens using e-cigarettes at school, it is critical to share resources to help them quit.

Resources for Teens

- Ask a trusted adult or friend for support
- Talk to a doctor about treatment options
- Visit smokefree.gov and teen.smokefree.gov/ quit-vaping
- Call 1-800-QUIT-NOW
- Text DITCHJUUL to 88709

References

- Wang TW, Neff LJ, Park-Lee E, Ren C, Cullen KA, King BA. E-cigarette Use Among Middle and High School Students United States, 2020. MMWR. 2020.
- 2. Centers for Disease Control and Prevention (CDC). Smoking & Tobacco. E-cigarettes shaped like USB flash drives: Information for parents, educators and health care providers. 2018.
- 3. Jackler RK et al. Tob Control (2019).
- 4. Rubinstein M et al. Pediatrics (2018).
- 5. Goniewicz ML et al. Tobacco Control. (2014)
- 6. Allen J, et al. Environ Health Perspect. (2016)
- 7. U.S. Department of Health and Human Services (USDHHS). A Report of the Surgeon General: How Tobacco Smoke Causes Disease. (2010)
- 8. U.S. Department of Health and Human Services (USDHHS). A Report of the Surgeon General: Preventing Tobacco Use among Youth and Young Adults. (2012)
- 9. U.S. Department of Health and Human Services (USDHHS). E-Cigarette Use Among Youth and Young Adults: A Report of the Surgeon General. (2016)
- 10. Williams M, et al. PLoS One. 2013
- 11. Olmedo P, et al. Environmental Health Perspectives (Online). 2018
- 12. Brook JS, Cohen P, Brook DW. (1998). Longitudinal study of co-occurring psychiatric disorders and substance use. J Am Acad Child Adolesc Psychiatry, 37, 322-30.
- 13. Counotte, D.S., et al. (2009). Long-Lasting Cognitive Deficits Resulting from Adolescent Nicotine Exposure in Rats. Neuropharmacology, 34, 299-306.
- 14. Mathers M, Toumbourou JW, Catalano RF, Williams J, Patton GC. (2006). Consequences of youth tobacco use: a review of prospective behavioural studies. Addiction, 101, 948-58.
- 15. Treur, J.L., Willemsen, G., Bartels, M., Geels, L.M., van Beek, J.H., Huppertz, C., et al. 2015. Smoking during adolescence as a risk factor for attention problems. Biol Psychiatry. 78(9): 656-663. 10.1016/j.biopsych.2014.06.019.
- 16. Marynak KL, Ali FRM, Schauer GL, et al. Tob Control 2019;28:685–688.